

CIRCULATION COPY
SUBJECT TO RECALL
IN TWO WEEKS

PREPRINT UCRL- 83370

Lawrence Livermore Laboratory

COLLISIONAL EFFECTS IN CARBON-13 ENRICHMENT BY CO₂ LASER

Jack B. Marling and Irving P. Herman

September, 1979

This paper was prepared for submission to CLEOS/ICF '80 in San Diego, California, February 26-28, 1980.

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.



DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

COLLISIONAL EFFECTS IN CARBON-13 ENRICHMENT BY CO₂ LASER[†]

Jack B. Marling and Irving P. Herman
University of California Lawrence Livermore Laboratory
P. O. Box 808, L-372
Livermore, California 94550
(415) 422-7293

ABSTRACT

Collisional effects have been evaluated in carbon-13 enrichment by multiple-photon dissociation of CF₃Br at 9.6 μ . Dissociation yields of both isotopic species increase with pressure. The single-step C-13 enrichment in C₂F₆ photo-product falls suddenly above 1 torr. Shortening the dissociation pulse to 2 nanoseconds permits operation to 225 torr.


[†] Work performed under the auspices of the U. S. Department of Energy by the Lawrence Livermore Laboratory under contract number W-7405-ENG-48.

COLLISIONAL EFFECTS IN CARBON-13 ENRICHMENT BY CO₂ LASER

Jack B. Marling and Irving P. Herman
University of California Lawrence Livermore Laboratory
P. O. Box 808, L-372
Livermore, California 94550
(415) 422-7293

SUMMARY

CO₂ laser multiple-photon dissociation of small halocarbon molecules has emerged as a very attractive route to enrichment of carbon-13.⁽¹⁾ Dissociation of CF₃Br near 9.6μ results in a single-step carbon-13 enrichment factor of 50-60, and is perhaps the best molecule for photochemical carbon-13 enrichment at room temperature.⁽¹⁾ We have examined the collisional effects in this process and have observed both an increase in photoproduct yield and a decrease in isotopic enrichment due to collisions. Under conditions of very low energy deposition, at wavelength of low molecular absorption coefficient, an increase in single-step enrichment occurs with increasing pressure, up to a critical pressure. Carbon-13 enrichment occurs via the steps



Both photoproduct yields and carbon-13 enrichment were determined from the C₂F₆ product. C₂F₆ product yields were determined by gas chromatography with TC detection, and carbon-13 enrichment determined from the mass 119, 120, 121 peaks of the C₂F₅ fragment in mass spectrometry.

At P(24), 9.6μ, the dissociation probability of ¹³CF₃Br was 10% below 0.1 torr pressure at an energy fluence of 26 J/cm². Collisions increased

the $^{13}\text{CF}_3\text{Br}$ dissociation probability monotonically to 100% as the total CF_3Br (containing 1.1% $^{13}\text{CF}_3\text{Br}$) was increased to 5 torr using a laser pulse duration of about 100 nanoseconds. However, the $^{12}\text{CF}_3\text{Br}$ dissociation yield also rose from 0.2% to 100% over the same pressure range. The net effect on the single-step enrichment factor is shown in the left-most curve of Figure 1. Figure 1 indicates a very sudden 40-fold drop in the C-13 enrichment factor as pressure doubles from 1 to 2 torr. We attribute this to a "thermal avalanche" effect. As pressure increases, absorption and hence energy loading within the focal volume of the laser beam increases. This local heating leads to population of higher J states in $^{12}\text{CF}_3\text{Br}$, which can now be excited by the laser beam leading to still further heating and increased absorption. The chosen wavelength of P(24), 9.6μ , is very much in the long-wavelength wing of $^{12}\text{CF}_3\text{Br}$ absorption, and very little $^{12}\text{CF}_3\text{Br}$ normally resides in these high J states. To verify this explanation, CF_3Br was dissociated at P(36), 9.7μ , where very little absorption occurs for either isotopic species. Under these conditions of weak absorption the "thermal avalanche" did not occur until 5 torr. In fact, the single-step C-13 enrichment factor increased from 40 at 0.2 torr to 65 at 2 torr. An increase in isotopic enrichment factor with pressure has also been observed in hexafluoroacetone.⁽²⁾ Above 4 torr the "thermal avalanche" effect resulted in an order-of-magnitude decrease in the C-13 enrichment factor with only 20% increase in CF_3Br pressure.

A dramatic improvement in the operating pressure limits in carbon-13 enrichment was achieved by shortening the CO_2 laser pulse duration to 2 nanoseconds (short-pulse experiments were performed at the Los Alamos Scientific Laboratory in collaboration with Scott J. Thomas). Under these irradiation

conditions at the same P(24) wavelength, no "thermal avalanche" effect was observed. Results are shown in Figure 1, right-hand curve, and indicate only a very slow decrease in carbon-13 enrichment out to 225 torr, the maximum pressure utilized. Additional details of carbon-13 enrichment will be discussed.

- (1) M. Drouin et al., Chem. Phys. Lett. 60, 16 (1978).
- (2) P. A. Hackett, M. Gauthier, C. Willis, and R. Pilon, J. Chem. Phys. 71, 546 (1979).

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

NOTICE

"This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately-owned rights."




FIGURE 1

RECEIVED
JAN 10 1971
NATIONAL BUREAU OF STANDARDS

CO₂ laser techniques can be applied very effectively in separation of light isotopes such as carbon-13. This figure shows that the single-step carbon-13 enrichment factor is about 50 at low pressure using CF₃Br. The CF₃Br dissociates by high power pulsed CO₂ laser (P(24), 9.6μ) to yield carbon-13 enriched ¹³CF₃ fragments that recombine to produce ¹³C₂F₆. The ¹³C₂F₆ may be then converted into the desired C-13 bearing products, or it may undergo further laser enrichment to yield nearly pure carbon-13. The sharp decrease in C-13 enrichment at higher pressure may be greatly reduced by using shorter duration CO₂ laser pulses, as shown in the curve to the right. Higher pressure operation greatly increases the throughput and greatly reduces the photochemical reactor size.